

# A CHARACTERIZATION OF DEPTH 2 SUBFACTORS OF $\text{II}_1$ FACTORS

Dmitri Nikshych\*      Leonid Vainerman†

October 15, 1998

## Abstract

We characterize finite index depth 2 inclusions of type  $\text{II}_1$  factors in terms of actions of weak Kac algebras and weak  $C^*$ -Hopf algebras. If  $N \subset M \subset M_1 \subset M_2 \subset \dots$  is the Jones tower constructed from such an inclusion  $N \subset M$ , then  $B = M' \cap M_2$  has a natural structure of a weak  $C^*$ -Hopf algebra and there is a minimal action of  $B$  on  $M_1$  such that  $M$  is the fixed point subalgebra of  $M_1$  and  $M_2$  is isomorphic to the crossed product of  $M_1$  and  $B$ . This extends the well-known results for irreducible depth 2 inclusions.

## 1 Introduction

Let  $N \subset M$  be a finite index depth 2 inclusion of type  $\text{II}_1$  factors and  $N \subset M \subset M_1 \subset M_2 \subset \dots$  the corresponding Jones tower. It was announced by A. Ocneanu and was proved in [18], [3], [10] that if  $N \subset M$  is irreducible, i.e., such that  $N' \cap M = \mathbb{C}$ , then  $B = M' \cap M_2$  has a natural structure of a finite-dimensional Kac algebra and there is a canonical outer action of  $B$  on  $M_1$  such that  $M = M_1^B$ , the fixed point subalgebra of  $M_1$  with respect to this action, and  $M_2$  is isomorphic to the crossed product  $M_1 \rtimes B$ . The outerness condition is equivalent to the relative commutant  $M_1' \cap M_1 \rtimes B$  being trivial (such actions are also called minimal). In the case of an infinite index a similar description in terms of multiplicative unitaries and quantum groups was obtained in [4].

---

\*UCLA, Department of Mathematics, 405 Hilgard Avenue, Los Angeles, CA 90095-1555; E-mail: nikshych@math.ucla.edu

†URA CNRS (Case 191), Université Pierre et Marie Curie, 4, place Jussieu, F-75252, Paris Cedex 05 France; E-mail: vainerma@math.jussieu.fr

In this work we extend the above result to (in general, reducible, i.e., such that  $\mathbb{C} \subset N' \cap M$ ) finite index depth 2 inclusions of type  $\text{II}_1$  factors. We replace usual Kac algebras (Hopf  $C^*$ -algebras) by weak Kac algebras [11] or weak  $C^*$ -Hopf algebras [2]. A weak Kac algebra is a special case of a weak  $C^*$ -Hopf algebra characterized by the property  $S^2 = \text{id}$ . Weak Kac algebras naturally arise in the situations when the index  $[M : N]$  is integer, e.g., when the inclusion is given by the crossed product with a usual Kac algebra. It was shown in [11] that the category of weak Kac algebras is equivalent to those of generalized Kac algebras of T. Yamanouchi [21] (another proof of that can be found in [14]) and of Kac bimodules (an algebraic version of Hopf bimodules of J.-M. Vallin [19]). The advantage of the language of weak Kac algebras and weak  $C^*$ -Hopf algebras is that their defining axioms are clearly self-dual, so it is easy to work with both weak Kac algebra (weak  $C^*$ -Hopf algebra) and its dual simultaneously.

Let us mention that a possibility of characterizing finite index depth 2 inclusions in terms of weak  $C^*$ -Hopf algebras was suggested in [13]. For an arbitrary (possibly infinite) index  $M$ . Enock and J.-M. Vallin have obtained a similar description in terms of pseudo-multiplicative unitaries [5].

The paper is organized as follows.

In Section 2 (Preliminaries) we briefly review, following [11], [2] and [13], the basic definitions and facts of the theory of weak Kac algebras and weak  $C^*$ -Hopf algebras, including their actions on von Neumann algebras.

Section 3 is devoted to establishing a non-degenerate duality between the finite dimensional  $C^*$ -algebras  $A = N' \cap M_1$  and  $B = M' \cap M_2$ , which gives a natural coalgebra structures on them.

In Sections 4 and 5 we investigate the relations between algebra and coalgebra structures on  $B$ , following the general strategy of Szymanski's reasoning [18] based on the above duality. It turns out that the square of the corresponding antipode is implemented by a positive invertible element determined by Index  $\tau|_{M' \cap M_1}$ , the Watatani index [20] of the restriction of the Markov trace  $\tau$  on  $M' \cap M_1$ . That is why it is natural to consider the cases of scalar and non-scalar Index  $\tau|_{M' \cap M_1}$  in which the antipode is respectively involutive and non-involutive. The main result here is that in the mentioned cases  $B$  and  $A$  are biconnected weak Kac algebras and weak  $C^*$ -Hopf algebras respectively (they are usual Kac algebras iff the inclusion  $N \subset M$  is irreducible). We also prove in Section 4, that if  $[M : N]$  is an integer which has no divisors of the form  $n^2$ ,  $n > 1$ , then the inclusion is irreducible and  $B$  is a Kac algebra acting outerly on  $M_1$ . In particular, if  $[M : N] = p$  is

prime, then  $B$  must be the group algebra of the cyclic group  $G = \mathbb{Z}/p\mathbb{Z}$ .

Finally, in Section 6 we show that there exists a canonical (left) minimal action of  $B$  on  $M_1$  such that  $M$  is the fixed point subalgebra of  $M_1$  with respect to this action, and  $M_2$  is isomorphic to  $M_1 \rtimes B$ , the crossed product of  $M_1$  and  $B$ . The minimality condition means that the relative commutant  $M'_1 \cap M_1 \rtimes B$  is minimal possible, in which case it is isomorphic to the Cartan subalgebra  $B_s \subset B$ .

It is important to stress that in the above situation one can take

$$\begin{array}{ccc} B^* & \subset & B^* \rtimes B \\ \cup & & \cup \\ B^* \cap B & \subset & B, \end{array}$$

where  $B^* = A$ , as a canonical commuting square [17] of the inclusion  $M_1 \subset M_2$ . The above square, and thus the equivalence class of inclusions, is completely determined by  $B$ . This implies that every biconnected weak  $C^*$ -Hopf algebra has at most one minimal action on a given  $\text{II}_1$  factor and thus correspond to no more than one (up to equivalence) finite index depth 2 subfactor. Note that any biconnected weak Kac algebra admits a unique minimal action on the hyperfinite  $\text{II}_1$  factor [12].

Let us remark that this characterization of depth 2 inclusions means that weak Kac algebras provide a good setting for studying actions of usual Kac algebras on  $\text{II}_1$  factors, since any (not necessarily minimal) action of a Kac algebra produces a depth 2 inclusion and one can canonically associate with this action a weak Kac algebra completely describing it. More details on this will be published elsewhere.

**Acknowledgements.** The first author would like to thank S. Popa and E. Effros for helpful advices, and E. Vaysleb for his useful comments on the preliminary version of this paper. The second author is deeply grateful to M. Enock and J.-M. Vallin for many important discussions.

## 2 Preliminaries

Our main references to finite dimensional weak  $C^*$ -Hopf algebras are [2] and [14]. Weak Kac algebras, a special case of this notion characterized by the property  $S^2 = \text{id}$ , were considered in [11]. These objects generalize both finite groupoid algebras and usual Kac algebras.

A *weak Kac algebra*  $B$  is a finite dimensional  $C^*$ -algebra equipped with the *comultiplication*  $\Delta : B \rightarrow B \otimes B$ , *counit*  $\varepsilon : B \rightarrow \mathbb{C}$ , and *antipode*

$S : B \rightarrow B$ , such that  $(\Delta, \varepsilon)$  defines a coalgebra structure on  $B$  and the following axioms hold for all  $b, c \in B$  (we use Sweedler's notation  $\Delta(b) = b_{(1)} \otimes b_{(2)}$  for the comultiplication) :

- (1)  $\Delta$  is a  $*$ -preserving (but not necessarily unital) homomorphism :

$$\Delta(bc) = \Delta(b)\Delta(c), \quad \Delta(b^*) = \Delta(b)^{* \otimes *},$$

- (2) The target counital map  $\varepsilon^t$ , defined by  $\varepsilon^t(b) = \varepsilon(1_{(1)}b)1_{(2)}$ , satisfies the relations

$$b\varepsilon^t(c) = \varepsilon(b_{(1)}c)b_{(2)}, \quad b_{(1)} \otimes \varepsilon^t(b_{(2)}) = 1_{(1)}b \otimes 1_{(2)},$$

- (3)  $S$  is an anti-algebra and anti-coalgebra map such that  $S^2 = \text{id}$ ,  $(S \circ *) = (* \circ S)$ , and

$$b_{(1)}S(b_{(2)}) = \varepsilon^t(b).$$

If instead of the conditions  $S^2 = \text{id}$  and  $(S \circ *) = (* \circ S)$  we have a less restrictive property  $(S \circ *)^2 = \text{id}$ , then  $B$  is called a *weak  $C^*$ -Hopf algebra*.

Note that the axioms (2) and (3) above are equivalent to the following axioms for the source counital map  $\varepsilon^s(b) = 1_{(1)}\varepsilon(b1_{(2)})$  :

$$(2') \quad \varepsilon^s(c)b = b_{(1)}\varepsilon(cb_{(2)}), \quad \varepsilon^s(b_{(1)}) \otimes b_{(2)} = 1_{(1)} \otimes b1_{(2)},$$

$$(3') \quad S(b_{(1)})b_{(2)} = \varepsilon^s(b).$$

The dual vector space  $B^*$  has a natural structure of a weak Kac algebra (weak  $C^*$ -Hopf algebra) given by dualizing the structure operations of  $B$ , see [2], [11].

The main difference between weak Kac ( $C^*$ -Hopf) algebras and classical Kac algebras is that the images of the counital maps are, in general, non-trivial unital  $C^*$ -subalgebras of  $B$ , called *Cartan subalgebras* (note that we have  $\varepsilon^t \circ \varepsilon^t = \varepsilon^t$  and  $\varepsilon^s \circ \varepsilon^s = \varepsilon^s$ ) :

$$\begin{aligned} B_t &= \{x \in B \mid \varepsilon^t(x) = x\} = \{x \in B \mid \Delta(x) = x1_{(1)} \otimes 1_{(2)} = 1_{(1)}x \otimes 1_{(2)}\}, \\ B_s &= \{x \in B \mid \varepsilon^s(x) = x\} = \{x \in B \mid \Delta(x) = 1_{(1)} \otimes x1_{(2)} = 1_{(1)} \otimes 1_{(2)}x\}. \end{aligned}$$

The Cartan subalgebras commute :  $[B_t, B_s] = 0$ , also we have  $S \circ \varepsilon^s = \varepsilon^t \circ S$  and  $S(B_t) = B_s$ . We say that  $B$  is *connected* [12] if  $B_t \cap Z(B) = \mathbb{C}$  (where  $Z(B)$  denotes the center of  $B$ ), i.e., if the inclusion  $B_t \subset B$  is connected.  $B$  is connected iff  $B_t^* \cap B_s^* = \mathbb{C}$  ([12], Proposition 3.11). We say that  $B$  is *biconnected* if both  $B$  and  $B^*$  are connected.

Weak Kac ( $C^*$ -Hopf) algebras have integrals in the following sense.

There exists a unique projection  $p \in B$ , called a *Haar projection*, such that for all  $x \in B$  :

$$xp = \varepsilon^t(x)p, \quad S(p) = p, \quad \varepsilon^t(p) = 1.$$

There exists a unique positive functional  $\phi$  on  $B$ , called a *normalized Haar functional* (which is a trace iff  $B$  is a weak Kac algebra), such that

$$(\text{id} \otimes \phi)\Delta = (\varepsilon^t \otimes \phi)\Delta, \quad \phi \circ S = S, \quad \phi \circ \varepsilon^t = \varepsilon.$$

The following notions of action, crossed product, and fixed point subalgebra were introduced in [13].

A (left) *action* of a weak Kac ( $C^*$ -Hopf algebra)  $B$  on a von Neumann algebra  $M$  is a linear map

$$B \otimes M \ni b \otimes x \mapsto (b \triangleright x) \in M$$

defining a structure of a left  $B$ -module on  $M$  such that for all  $b \in B$  the map  $b \otimes x \mapsto (b \triangleright x)$  is weakly continuous and

- (1)  $b \triangleright xy = (b_{(1)} \triangleright x)(b_{(2)} \triangleright y)$ ,
- (2)  $(b \triangleright x)^* = S(b)^* \triangleright x^*$ ,
- (3)  $b \triangleright 1 = \varepsilon^t(b) \triangleright 1$ , and  $b \triangleright 1 = 0$  iff  $\varepsilon^t(b) = 0$ .

A *crossed product* algebra  $M \rtimes B$  is constructed as follows. As a  $\mathbb{C}$ -vector space it is  $M \otimes_{B_t} B$ , where  $B$  is a left  $B_t$ -module via multiplication and  $M$  is a right  $B_t$ -module via multiplication by the image of  $B_t$  under  $z \mapsto (z \triangleright 1)$ ; that is, we identify

$$x(z \triangleright 1) \otimes b \equiv x \otimes zb$$

for all  $x \in M$ ,  $b \in B$ ,  $z \in B_t$ . Let  $[x \otimes b]$  denote the class of  $x \otimes b$ . A  $*$ -algebra structure on  $M \rtimes B$  is defined by

$$[x \otimes b][y \otimes c] = [x(b_{(1)} \triangleright y) \otimes b_{(2)}c], \quad [x \otimes b]^* = [(b_{(1)}^* \triangleright x^*) \otimes b_{(2)}^*]$$

for all  $x, y \in A$ ,  $b, c \in B$ . It is possible to show that this abstractly defined  $*$ -algebra  $M \rtimes B$  is  $*$ -isomorphic to a weakly closed algebra of operators on some Hilbert space [13], i.e.,  $M \rtimes B$  is a von Neumann algebra.

The collection  $M^B = \{x \in M \mid b \triangleright x = \varepsilon^t(b) \triangleright x, \forall b \in B\}$  is a von Neumann subalgebra of  $M$ , called a *fixed point subalgebra*.

The relative commutant  $M' \cap M \rtimes B$  always contains a  $*$ -subalgebra isomorphic to  $B_s$ . Indeed, if  $z \in B_s$ , then it follows easily from the axioms of a weak  $C^*$ -Hopf algebra that  $\Delta(z) = 1_{(1)} \otimes 1_{(2)} z$ , therefore

$$\begin{aligned} [1 \otimes z][x \otimes 1] &= [(z_{(1)} \triangleright x) \otimes z_{(2)}] = [(1_{(1)} \triangleright x) \otimes 1_{(2)} z] \\ &= [x \otimes z] = [x \otimes 1][1 \otimes z], \end{aligned}$$

for all  $x \in M$ , and  $B_s \subset M' \cap M \rtimes B$ . We say that the action  $\triangleright$  is *minimal* if  $B_s = M' \cap M \rtimes B$ .

### 3 Duality between relative commutants

Let  $N \subset M$  be a depth 2 inclusion of type  $\text{II}_1$  factors with a finite index  $[M : N] = \lambda^{-1}$  and

$$N \subset M \subset M_1 \subset M_2 \subset \dots$$

be the corresponding Jones tower,  $M_1 = \langle M, e_1 \rangle$ ,  $M_2 = \langle M_1, e_2 \rangle, \dots$ , where  $e_1 \in N' \cap M_1$ ,  $e_2 \in M' \cap M_2, \dots$  are the Jones projections. The depth 2 condition means that  $N' \cap M_2$  is the basic construction of the inclusion  $N' \cap M \subset N' \cap M_1$ . Let  $\tau$  be the normalized (Markov) trace on  $M_2$ .

With respect to this trace, the square of algebras in the upper right corner of the diagram below

$$\begin{array}{ccccc} N' \cap M & \subset & N' \cap M_1 & \subset & N' \cap M_2 \\ & & \cup & & \cup \\ & & M' \cap M_1 & \subset & M' \cap M_2 \\ & & & & \cup \\ & & & & M'_1 \cap M_2. \end{array}$$

is commuting ( $E_{M_1} \circ E_{M'} = E_{M'} \circ E_{M_1}$  on  $N' \cap M_2$ ) and non-degenerate, i.e.,  $N' \cap M_2 = (N' \cap M_1)(M' \cap M_2)$ . This square is called a standard (or canonical) commuting square of the inclusion  $M_1 \subset M_2$  [17].

Let us denote

$$\begin{aligned} A &= N' \cap M_1, & B &= M' \cap M_2, \\ A_t &= N' \cap M, & A_s &= M' \cap M_1 = B_t, & B_s &= M'_1 \cap M_2. \end{aligned}$$

Note that  $A_t$  commutes with  $B$ ,  $B_s$  commutes with  $A$ , and  $A \cap B = A_s = B_t$ .

The next lemma will be frequently used in the sequel without specific reference.

**Lemma 3.1**  $(N' \cap M_2)e_2 = Ae_2$  and  $(N' \cap M_2)e_1 = Be_1$ . More precisely, for any  $x \in N' \cap M_2$  we have

$$xe_2 = \lambda^{-1}E_{M_1}(xe_2)e_2, \quad xe_1 = \lambda^{-1}E_{M'}(xe_1)e_1.$$

*Proof.* This statement is a special case of ([15], Lemma 1.2) since  $N' \cap M_2$  is the basic construction for the inclusions  $N' \cap M \subset N' \cap M_1$  and  $M'_1 \cap M_2 \subset M' \cap M_2$  with the corresponding Jones projections  $e_2$  and  $e_1$  respectively.

Let us denote  $d = \dim(M' \cap M_1)$ .

**Proposition 3.2** *The form*

$$\langle a, b \rangle = d\lambda^{-2}\tau(ae_2e_1b), \quad a \in A, b \in B$$

*defines a non-degenerate duality between A and B.*

*Proof.* If  $a \in A$  is such that  $\langle a, B \rangle = 0$ , then

$$\tau(ae_2e_1B) = \tau(ae_2e_1(N' \cap M_2)) = 0,$$

therefore, using the Markov property of  $\tau$  and properties of Jones projections, we get

$$\tau(aa^*) = \lambda^{-1}\tau(ae_2a^*) = \lambda^{-2}\tau(ae_2e_1(e_2a^*)) = 0,$$

so  $a = 0$ . Similarly for  $b \in B$ .

**Definition 3.3** Using the form  $\langle, \rangle$  define the comultiplication  $\Delta_B$ , counit  $\varepsilon_B$ , and antipode  $S_B$  as follows :

$$\begin{aligned} \Delta_B : B &\rightarrow B \otimes B : & \langle a_1a_2, b \rangle &= \langle a_1, b_{(1)} \rangle \langle a_2, b_{(2)} \rangle, \\ \varepsilon_B : B &\rightarrow \mathbb{C} : & \varepsilon_B(b) &= \langle 1, b \rangle = \lambda^{-1}d\tau(be_2), \\ S_B : B &\rightarrow B : & \langle a, S_B(b) \rangle &= \overline{\langle a^*, b^* \rangle}, \end{aligned}$$

for all  $a, a_1, a_2 \in A$  and  $b \in B$ . Similarly, we define  $\Delta_A$ ,  $\varepsilon_A$ , and  $S_A$ .

Clearly,  $(B, \Delta_B, \varepsilon_B)$  (resp.  $(A, \Delta_A, \varepsilon_A)$ ) becomes a coalgebra. Let us investigate the relations between the algebra and coalgebra structures on  $B$ .

## 4 Weak Kac algebra structure on $M' \cap M_2$ (the case of a scalar Watatani index of $\tau|_{M' \cap M_1}$ )

**Lemma 4.1** *For all  $a \in A$  and  $b_1, b_2 \in B$  we have*

$$\langle a, b_1 b_2 \rangle = \lambda^{-1} \langle E_{M_1}(b_2 a e_2), b_1 \rangle.$$

*Proof.* Using the definition of  $\langle, \rangle$  we have

$$\begin{aligned} \langle a, b_1 b_2 \rangle &= d\lambda^{-2} \tau(b_2 a e_2 e_1 b_1) = d\lambda^{-3} \tau(E_{M_1}(b_2 a e_2) e_2 e_1 b_1) \\ &= \lambda^{-1} \langle E_{M_1}(b_2 a e_2), b_1 \rangle. \end{aligned}$$

**Proposition 4.2** *Let  $\varepsilon_B^t(b) = \varepsilon_B(1_{(1)} b) 1_{(2)}$ . Then  $\varepsilon_B^t(b) = \lambda^{-1} E_{M_1}(b e_2)$  and*

$$\langle a, \varepsilon_B^t(b) \rangle = d\lambda^{-2} \tau(a e_1 b e_2) = \lambda^{-1} \langle E_M(a e_1), b \rangle.$$

*Proof.* Using Lemma 4.1, definitions of  $\Delta_B$  and  $\varepsilon_B$ , we have

$$\begin{aligned} \langle a, \varepsilon_B(1_{(1)} b) 1_{(2)} \rangle &= \langle 1, 1_{(1)} b \rangle \langle a, 1_{(2)} \rangle \\ &= \langle \lambda^{-1} E_{M_1}(b e_2), 1_{(1)} \rangle \langle a, 1_{(2)} \rangle \\ &= \langle \lambda^{-1} E_{M_1}(b e_2) a, 1 \rangle = \langle a, \lambda^{-1} E_{M_1}(b e_2) \rangle, \end{aligned}$$

from where the first statement follows. For the second one, we have, using the  $\lambda$ -Markov property and the fact that  $e_2$  commutes with  $M$ ,

$$\begin{aligned} \langle a, \lambda^{-1} E_{M_1}(b e_2) \rangle &= d\lambda^{-2} \tau(a e_1 e_2 \lambda^{-1} E_{M_1}(b e_2)) \\ &= d\lambda^{-2} \tau(a e_1 \lambda^{-1} E_{M_1}(b e_2) e_2) \\ &= d\lambda^{-2} \tau(a e_1 b e_2) = d\lambda^{-3} \tau(E_M(a e_1) e_1 b e_2) \\ &= d\lambda^{-3} \tau(E_M(a e_1) e_2 e_1 b) = \lambda^{-1} \langle E_M(a e_1), b \rangle. \end{aligned}$$

**Proposition 4.3** *For all  $b, c \in B$  we have*

$$b_{(1)} \otimes \varepsilon_B^t(b_{(2)}) = 1_{(1)} b \otimes 1_{(2)}, \quad b \varepsilon_B^t(c) = \varepsilon_B(b_{(1)} c) b_{(2)}.$$



*Proof.* For all  $a_1, a_2 \in A$  we compute, using Lemma 4.1 and Proposition 4.2 :

$$\begin{aligned}
\langle a_1, b_{(1)} \rangle \langle a_2, \varepsilon_B^t(b_{(2)}) \rangle &= \langle a_1 \lambda^{-1} E_M(a_2 e_1), b \rangle \\
&= \lambda^{-2} \langle E_{M_1}(ba_1 E_M(a_2 e_1) e_2), 1 \rangle \\
&= \lambda^{-2} \langle E_{M_1}(ba_1 e_2) E_M(a_2 e_1), 1 \rangle \\
&= d \lambda^{-3} \tau(E_{M_1}(ba_1 e_2) E_M(a_2 e_1) e_1) \\
&= d \lambda^{-2} \tau(E_{M_1}(ba_1 e_2) a_2 e_1) \\
&= \lambda^{-1} \langle E_{M_1}(ba_1 e_2) a_2, 1 \rangle \\
&= \langle \lambda^{-1} E_{M_1}(ba_1 e_2), 1_{(1)} \rangle \langle a_2, 1_{(2)} \rangle \\
&= \langle a_1, 1_{(1)} b \rangle \langle a_2, 1_{(2)} \rangle, \\
\langle a, b \varepsilon_B^t(c) \rangle &= \langle \varepsilon_B^t(c) a, b \rangle \\
&= \langle \lambda^{-1} E_{M_1}(ce_2) a, b \rangle \\
&= \langle \lambda^{-1} E_{M_1}(ce_2), b_{(1)} \rangle \langle a, b_{(2)} \rangle \\
&= \langle 1, b_{(1)} c \rangle \langle a, b_{(2)} \rangle \\
&= \langle a, \varepsilon_B(b_{(1)} c) b_{(2)} \rangle.
\end{aligned}$$

Since the duality is non-degenerate, the result follows.

The antipode map assigns to each  $b \in B$  a unique element  $S_B(b) \in B$  such that  $\tau(ae_2e_1S_B(b)) = \tau(be_1e_2a)$  for all  $a \in A$ , or, equivalently,

$$E_{M_1}(be_1e_2) = E_{M_1}(e_2e_1S_B(b)).$$

Taking  $a = e_1$  and using the  $\lambda$ -Markov property of  $e_1$  we get  $\tau \circ S_B = \tau$ . Similarly,  $E_{M'}(S_A(a)e_2e_1) = E_{M'}(e_1e_2a)$  and  $\tau \circ S_A = \tau$ .

**Remark 4.4** Note that the condition  $E_{M_1}(be_1e_2) = E_{M_1}(e_2e_1S_B(b))$  implies that

$$E_{M_1}(bx e_2) = E_{M_1}(e_2x S_B(b)) \quad \text{for all } x \in M_1.$$

Indeed, any  $x \in M_1$  can be written as  $x = \sum x_i e_1 y_i$  with  $x_i, y_i \in M \subset B'$ . Similarly, we have

$$E_{M'}(S_A(a) y e_1) = E_{M'}(e_1 y a) \quad \text{for all } y \in M'.$$

**Proposition 4.5** *The following identities hold :*

$$(i) \ S_B(b) = \lambda^{-3} E_{M'}(e_1 e_2 E_{M_1}(b e_1 e_2)),$$

$$(ii) \ S_B(B_s) = B_t,$$

$$(iii) \ S_B^2(b) = b \text{ and } S_B(b)^* = S_B(b^*),$$

$$(iv) \ S_B(bc) = S_B(c)S_B(b) \text{ and } \Delta_B(S_B(b)) = \varsigma(S_B \otimes S_B)\Delta_B(b).$$

*Proof.* (i) We have

$$\begin{aligned} S_B(b) &= \lambda^{-1} E_{M'}(e_1 S_B(b)) = \lambda^{-2} E_{M'}(e_1 e_2 e_1 S_B(b)) \\ &= \lambda^{-3} E_{M'}(e_1 e_2 E_{M_1}(e_2 e_1 S_B(b))) = \lambda^{-3} E_{M'}(e_1 e_2 E_{M_1}(b e_1 e_2)). \end{aligned}$$

(ii) If  $z \in B_s$  then  $z e_2 = e_2 z$  and by the explicit formula (i) we get,

$$\begin{aligned} S_B(z) &= \lambda^{-3} E_{M'}(e_1 e_2 E_{M_1}(e_1 z e_2)) = \lambda^{-2} E_{M'}(e_1 E_{M_1}(z e_2)) \\ &= \lambda^{-1} E_{M_1}(z e_2) = \varepsilon_B^t(z) \in B_t. \end{aligned}$$

(iii) Since  $E_{M_1}$  preserves  $*$ , we get  $E_{M_1}(e_2 e_1 b^*) = E_{M_1}(S_B(b)^* e_2 e_1)$ , from where  $S_B(S_B(b)^*)^* = b$ . Next, using Lemma 4.1, Remark 4.4, and the  $\lambda$ -Markov property of  $e_2$ , we compute

$$\begin{aligned} \tau(a e_2 e_1 b) &= \lambda^{-1} \tau(E_{M_1}(b a e_2) e_2 e_1) = \lambda^{-1} \tau(E_{M_1}(e_2 a S_B(b)) e_2 e_1) \\ &= \lambda^{-1} \tau(e_2 E_{M_1}(e_2 a S_B(b)) e_1) = \tau(e_2 a S_B(b) e_1) \\ &= \tau(S_B(b) e_1 e_2 a) = \tau(a e_2 e_1 S_B^2(b)). \end{aligned}$$

therefore,  $S_B^2(b) = b$  and  $S_B(b)^* = S_B(b^*)$ .

(iv) Using Remark 4.4, we have

$$\begin{aligned} \tau(a e_2 e_1 S_B(bc)) &= \tau(b c e_1 e_2 a) = \lambda^{-1} \tau(c e_1 e_2 E_{M_1}(e_2 a b)) \\ &= \lambda^{-1} \tau(E_{M_1}(e_2 a b) e_2 e_1 S_B(c)) \\ &= \lambda^{-1} \tau(E_{M_1}(S_B(b) a e_2) e_2 e_1 S_B(c)) \\ &= \tau(a e_2 e_1 S_B(c) S_B(b)), \end{aligned}$$

which proves that  $\langle a, S_B(bc) \rangle = \langle a, S_B(c) S_B(b) \rangle$ . Similarly, one can prove that  $S_A$  is anti-multiplicative, and since  $\langle a, S_B(b) \rangle = \langle S_A(a), b \rangle$ , the second part of (iv) follows.

Let  $\{f_{kl}^\alpha\}$  be a system of matrix units in  $B_t = M' \cap M_1 = \oplus_\alpha M_{m_\alpha}(\mathbb{C})$ , where  $\sum m_\alpha^2 = d$ , and let  $\tau_\alpha = \tau(f_{kk}^\alpha)$ .

**Proposition 4.6** *The explicit formula for  $\Delta_B(1)$  is*

$$\Delta_B(1) = \sum_{\alpha kl} \frac{1}{d\tau_\alpha} S_B(f_{kl}^\alpha) \otimes f_{lk}^\alpha.$$

*In particular,  $\Delta_B(1)$  is a positive element in  $B_s \otimes B_t$ .*

*Proof.* Note that the map  $x \mapsto \sum_{\alpha kl} \frac{\tau(x f_{lk}^\alpha)}{\tau_\alpha} f_{kl}^\alpha$  defines the  $\tau$ -preserving conditional expectation on  $B_t$ . For all  $a_1, a_2 \in A$  we have

$$\begin{aligned} \sum_{\alpha kl} \frac{1}{d\tau_\alpha} \langle a_1, S_B(f_{kl}^\alpha) \rangle \langle a_2, f_{lk}^\alpha \rangle &= \\ &= d^2 \lambda^{-4} \sum_{\alpha kl} \frac{1}{d\tau_\alpha} \tau(a_1 e_2 e_1 S_B(f_{kl}^\alpha)) \tau(a_2 e_2 e_1 f_{lk}^\alpha) \\ &= d \lambda^{-3} \sum_{\alpha kl} \tau(f_{kl}^\alpha e_1 e_2 a_1) \frac{\tau(a_2 e_1 f_{lk}^\alpha)}{\tau_\alpha} \\ &= d \lambda^{-3} \tau(E_{M'}(a_2 e_1) e_1 e_2 a_1) \\ &= d \lambda^{-2} \tau(a_1 a_2 e_1 e_2) = \langle a_1 a_2, 1 \rangle, \end{aligned}$$

which proves the statement.

**Corollary 4.7**  $\Delta_B(1) = \sum_{\alpha kl} \frac{1}{m_\alpha} S_B(f_{kl}^\alpha) \otimes f_{lk}^\alpha H$ , where  $H$  is canonically defined by

$$H = S_B(1_{(1)}) 1_{(2)} = \frac{1}{d} \sum_{\alpha} \frac{m_\alpha}{\tau_\alpha} \sum_k f_{kk}^\alpha = \frac{1}{d} \text{Index} \tau|_{M' \cap M_1} \in Z(B_t),$$

where  $\text{Index} \tau|_{M' \cap M_1}$  is the Watatani index [20] of the restriction of  $\tau$  to  $M' \cap M_1$  and  $Z(\cdot)$  denotes the center of the algebra. We also have  $\tau(H) = 1$ .

**Proposition 4.8** *For all  $b \in B$  we have  $\varepsilon_B^t(b_{(1)}) b_{(2)} = Hb$ .*

*Proof.* Applying  $E_{M'}$  to both sides of  $E_{M_1}(b^* e_1 e_2) = E_{M_1}(e_2 e_1 S_B(b^*))$  and using the relation  $E_{M_1} \circ E_{M'} = E_{M'} \circ E_{M_1}$ , we get

$$E_{M_1}(b^* e_2) = E_{M_1}(e_2 S_B(b^*))$$

which means that  $\varepsilon_B^t(b^*) = \varepsilon_B^t(S_B(b))^*$ . Using Propositions 4.3, 4.5(iv), and Corollary 4.7 we get  $S_B(b_{(1)}) \varepsilon_B^t(b_{(2)}) = S_B(b)H$ , from where  $HS_B(b^*) = \varepsilon_B^t(b_{(2)})^* S(b_{(1)}^*)$ . Replacing  $S_B(b^*)$  by  $b$ , we get the result.

Let  $\{s_{jk}^\alpha\}$  be a basis consisting of matrix units of  $A$  and  $\{v_{jk}^\alpha\}$  be a basis of comatrix units of  $B$  dual to each other, i.e.,

$$\langle v_{jk}^\alpha, s_{pq}^\beta \rangle = \delta_{\alpha\beta} \delta_{jp} \delta_{kq}.$$

We have  $\Delta_B(v_{jk}^\alpha) = \sum_l v_{jl}^\alpha \otimes v_{lk}^\alpha$  and  $\varepsilon_B(v_{jk}^\alpha) = \delta_{jk}$ .

**Lemma 4.9** *Let  $/\alpha/ = \tau(s_{kk}^\alpha)$ . The following identities hold true :*

- (i)  $E_{M_1}(e_2 e_1 v_{jk}^\alpha) = d^{-1} \lambda^2 / \alpha /^{-1} s_{kj}^\alpha$ ,
- (ii)  $E_{M_1}(v_{jk}^\alpha e_1 e_2) = d^{-1} \lambda^2 / \alpha /^{-1} S_A(s_{kj}^\alpha)$ ,
- (iii)  $\lambda^{-1} E_{M'}(S_A(s_{pq}^\beta) v_{ij}^\alpha e_1) = \delta_{\alpha\beta} \delta_{ip} v_{qj}^\alpha$ ,
- (iv)  $S_B(v_{jk}^\alpha) = (v_{kj}^\alpha)^*$ .

*Proof.* (i) We can directly compute :

$$\begin{aligned} d\lambda^{-2} / \alpha / \tau(s_{pq}^\beta E_{M_1}(e_2 e_1 v_{jk}^\alpha)) &= / \alpha / \langle s_{pq}^\beta, v_{jk}^\alpha \rangle \\ &= / \alpha / \delta_{\alpha\beta} \delta_{jp} \delta_{kq} \\ &= \tau(s_{kj}^\alpha s_{pq}^\beta), \end{aligned}$$

therefore, we have  $E_{M_1}(e_2 e_1 v_{jk}^\alpha) = d^{-1} \lambda^2 / \alpha /^{-1} s_{kj}^\alpha$  by the faithfulness of  $\tau$ .

(ii) Similarly to (i), we compute

$$\begin{aligned} d\lambda^{-2} / \alpha / \tau(E_{M_1}(v_{jk}^\alpha e_1 e_2) S_A(s_{pq}^\beta)) &= / \alpha / \langle s_{pq}^\beta, v_{jk}^\alpha \rangle \\ &= / \alpha / \delta_{\alpha\beta} \delta_{jp} \delta_{kq} \\ &= \tau(S_A(s_{kj}^\alpha) S_A(s_{pq}^\beta)), \end{aligned}$$

and since  $\tau \circ S_A = \tau$ , the result follows.

(iii) Using Remark 4.4, we have

$$\begin{aligned} \langle s_{rt}^\gamma, \lambda^{-1} E_{M'}(S_A(s_{pq}^\beta) v_{ij}^\alpha e_1) \rangle &= \langle s_{rt}^\gamma, \lambda^{-1} E_{M'}(e_1 v_{ij}^\alpha s_{pq}^\beta) \rangle \\ &= \langle s_{pq}^\beta s_{rt}^\gamma, v_{ij}^\alpha \rangle \\ &= \delta_{\alpha\gamma} \delta_{qr} \delta_{ip} \delta_{\alpha\beta} \delta_{tj} \\ &= \delta_{\alpha\beta} \delta_{ip} \langle s_{rt}^\gamma, v_{qj}^\alpha \rangle. \end{aligned}$$

(iv) Using part (i), we have

$$\begin{aligned} E_{M_1}((v_{kj}^\alpha)^* e_1 e_2) &= E_{M_1}(e_2 e_1 v_{kj}^\alpha)^* = d^{-1} \lambda^2 / \alpha /^{-1} s_{kj}^\alpha \\ &= E_{M_1}(e_2 e_1 v_{jk}^\alpha) = E_{M_1}(S_B(v_{jk}^\alpha) e_1 e_2), \end{aligned}$$

and the result follows from the injectivity of the map  $b \mapsto E_{M_1}(b e_1 e_2)$ .

**Corollary 4.10**  $\Delta_B(b^*) = \Delta_B(b)^{* \otimes *}$ , i.e.,  $\Delta_B$  is a  $*$ -preserving map.

*Proof.* Using Lemmas 4.9(iv) and Lemmas 4.5(iv), we have

$$\begin{aligned}\Delta_B((v_{jk}^\alpha)^*) &= \Delta_B(S_B(v_{kj}^\alpha)) = \Sigma_i S_B(v_{ij}^\alpha) \otimes S_B(v_{ki}^\alpha) \\ &= \Sigma_i (v_{ji}^\alpha)^* \otimes (v_{ik}^\alpha)^* = \Delta_B(v_{jk}^\alpha)^{* \otimes *}.\end{aligned}$$

**Proposition 4.11**  $v_{ij}^\alpha e_1 = \lambda^{-1} \sum_k E_{M_1}(v_{ik}^\alpha e_1 e_2) H^{-1} v_{kj}^\alpha$ .

*Proof.* By Lemma 4.9(ii), all we need to show is

$$v_{ij}^\alpha e_1 = d^{-1} \lambda / \alpha /^{-1} \sum_k S_A(s_{ki}^\alpha) H^{-1} v_{kj}^\alpha.$$

Since  $N' \cap M_2$  is spanned by the elements of the form  $v_{rt}^\gamma S_A(s_{pq}^\beta)$ , it suffices to verify that

$$\tau(v_{rt}^\gamma S_A(s_{pq}^\beta) v_{ij}^\alpha e_1) = d^{-1} \lambda / \alpha /^{-1} \sum_k \tau(v_{rt}^\gamma S_A(s_{pq}^\beta) S_A(s_{ki}^\alpha) H^{-1} v_{kj}^\alpha),$$

or, equivalently,

$$E_{M'}(S_A(s_{pq}^\beta) v_{ij}^\alpha e_1) = \delta_{\alpha\beta} \delta_{ip} \lambda d^{-1} / \alpha /^{-1} \sum_k E_{M'}(S_A(s_{kq}^\beta) H^{-1} v_{kj}^\alpha).$$

Using Lemma 4.9(iii), we can reduce the proof to the verification of the relation

$$v_{qj}^\alpha = d^{-1} / \alpha /^{-1} \sum_k E_{M'}(S_A(s_{kq}^\alpha)) H^{-1} v_{kj}^\alpha.$$

By Lemma 4.9(ii),

$$E_{M'}(S_A(s_{kq}^\alpha)) = d \lambda^{-2} / \alpha / E_{M'} \circ E_{M_1}(v_{qk}^\alpha e_1 e_2) = d \lambda^{-1} / \alpha / E_{M_1}(v_{qk}^\alpha e_2),$$

therefore the previous relation is equivalent to

$$v_{qj}^\alpha = \lambda^{-1} \sum_k E_{M_1}(v_{qk}^\alpha e_2) H^{-1} v_{kj}^\alpha.$$

Since  $H \in Z(B_t)$ , this is precisely Proposition 4.8 with  $b = v_{qj}^\alpha$ , so the proof is complete.

**Corollary 4.12**  $bx = \lambda^{-1}E_{M_1}(b_{(1)}xe_2)H^{-1}b_{(2)}$  for all  $b \in B$  and  $x \in M_1$ .

*Proof.* Proposition 4.11 implies that  $be_1 = \lambda^{-1}E_{M_1}(b_{(1)}e_1e_2)H^{-1}b_{(2)}$  for all  $b \in B$ . As in Remark 4.4, any  $x \in M_1$  can be written as a finite sum  $x = \sum x_ie_2y_i$  with  $x_i, y_i \in M \subset B'$ , therefore, we have

$$\begin{aligned} bx &= \sum x_ib e_1 y_i = \sum x_i \lambda^{-1} E_{M_1}(b_{(1)}e_1e_2)H^{-1}b_{(2)}y_i \\ &= \lambda^{-1}E_{M_1}(b_{(1)} \sum x_ie_1y_ie_2)H^{-1}b_{(2)} \\ &= \lambda^{-1}E_{M_1}(b_{(1)}xe_2)H^{-1}b_{(2)}. \end{aligned}$$

**Proposition 4.13** For all  $x, y \in M_1$ ,

$$E_{M_1}(bx ye_2) = \lambda^{-1}E_{M_1}(b_{(1)}xe_2)H^{-1}E_{M_1}(b_{(2)}ye_2).$$

*Proof.* Multiplying the formula from Corollary 4.12 on the right by  $ye_2t$  with  $y, t \in M_1$  and taking  $\tau$  from both sides we get

$$\tau(bx ye_2t) = \lambda^{-1}\tau(E_{M_1}(b_{(1)}xe_2)H^{-1}b_{(2)}ye_2t),$$

for all  $t \in M_1$ , from where the result follows.

**Proposition 4.14**  $\Delta_B(bc) = \Delta_B(b)(1 \otimes H^{-1})\Delta_B(c)$ , for all  $b, c \in B$ .

*Proof.* By Lemma 4.1 and Proposition 4.13 we have for all  $a_1, a_2 \in A$  :

$$\begin{aligned} \langle a_1a_2, bc \rangle &= \\ &= \langle \lambda^{-1}E_{M_1}(ca_1a_2e_2), b \rangle \\ &= \langle \lambda^{-2}E_{M_1}(c_{(1)}a_1e_2)H^{-1}E_{M_1}(c_{(2)}a_2e_2), b \rangle \\ &= \langle \lambda^{-1}E_{M_1}(c_{(1)}a_1e_2), b_{(1)} \rangle \langle \lambda^{-1}E_{M_1}(H^{-1}c_{(2)}a_2e_2), b_{(2)} \rangle \\ &= \langle a_1, b_{(1)}c_{(1)} \rangle \langle a_2, b_{(2)}H^{-1}c_{(2)} \rangle, \end{aligned}$$

from where  $\Delta_B(bc) = b_{(1)}c_{(1)} \otimes b_{(2)}H^{-1}c_{(2)}$  which is the result.

**Proposition 4.15**  $b_{(1)}S_B(b_{(2)}H^{-1}) = \varepsilon_B^t(b)$ .

*Proof.* Using Corollary 4.10, Proposition 4.13 and Proposition 4.2 we have

$$\begin{aligned}
\langle a, b_{(1)}S_B(b_{(2)}H^{-1}) \rangle &= d\lambda^{-3}\tau(E_{M_1}(S_B(b_{(2)}H^{-1})ae_2)e_2e_1b_{(1)}) \\
&= d\lambda^{-3}\tau(E_{M_1}(e_2ab_{(2)}H^{-1})e_2e_1b_{(1)}) \\
&= d\lambda^{-3}\tau(E_{M_1}(e_2ab_{(2)}H^{-1})E_{M_1}(e_2e_1b_{(1)})) \\
&= d\lambda^{-2}\tau(E_{M_1}(e_2ae_1b)) \\
&= \langle a, \varepsilon_B^t(b) \rangle.
\end{aligned}$$

The next Corollary summarizes the properties of  $\Delta_B$ ,  $\varepsilon_B$  and  $S_B$ .

**Corollary 4.16**  $(\Delta_B, \varepsilon_B)$  defines a coalgebra structure on  $B$  such that

$$\Delta_B(bc) = \Delta_B(b)(1 \otimes H^{-1})\Delta_B(c) \quad \Delta_B(b^*) = \Delta_B(b)^{* \otimes *},$$

the map  $\varepsilon_B^t$ , defined by  $\varepsilon_B^t(b) = \varepsilon_B(1_{(1)}b)1_{(2)}$ , satisfies the relations

$$b_{(1)} \otimes \varepsilon_B^t(b_{(2)}) = 1_{(1)}b \otimes 1_{(2)}, \quad b\varepsilon_B^t(c) = \varepsilon_B(b_{(1)}c)b_{(2)},$$

and there is a  $*$ -preserving anti-algebra and anti-coalgebra involution  $S_B$  such that

$$b_{(1)}S_B(b_{(2)}H^{-1}) = \varepsilon_B^t(b),$$

for all  $b, c \in B$ .

**Theorem 4.17** The following conditions are equivalent :

- (i)  $(B, \Delta_B, \varepsilon_B, S_B)$  is a weak Kac algebra with the Haar projection  $e_2$  and the normalized Haar trace  $\phi(b) = d\tau(b)$ ,  $b \in B$ ,
- (ii)  $H = 1$ .

Moreover, if these conditions are satisfied, then  $\lambda^{-1}$  is an integer.

*Proof.* (i) $\Rightarrow$ (ii). If  $\Delta_B$  is an algebra homomorphism, then we must have  $\Delta_B(1) = \Delta_B(1)(1 \otimes H^{-1})$ , and applying  $(\varepsilon_B \otimes \text{id})$  we get  $H^{-1} = 1$ .

(ii) $\Rightarrow$ (i). Clearly, if  $H = 1$ , then  $(B, \Delta_B, \varepsilon_B, S_B)$  is a weak Kac algebra. For all  $b \in B$  we have, by Proposition 4.2 :

$$be_2 = \lambda^{-1}E_{M_1}(be_2)e_2 = \varepsilon_B^t(b)e_2,$$

and we easily get  $S_B(e_2) = e_2$  and  $\varepsilon_B^t(e_2) = 1$ , so  $e_2$  is the Haar projection in  $B$ .

Next, since  $\tau(b) = d^{-1}\langle e_1, b \rangle$ , we have by Proposition 4.2 :

$$\begin{aligned} \langle a, \varepsilon_B^t(b_{(1)})\tau(b_{(2)}) \rangle &= d^{-1}\langle a, \varepsilon_B^t(b_{(1)}) \rangle \langle e_1, b_{(2)} \rangle \\ &= d^{-1}\langle \lambda^{-1}E_M(ae_1), b_{(1)} \rangle \langle e_1, b_{(2)} \rangle \\ &= d^{-1}\langle \lambda^{-1}E_M(ae_1)e_1, b \rangle \\ &= d^{-1}\langle ae_1, b \rangle = \langle a, b_{(1)}\tau(b_{(2)}) \rangle, \end{aligned}$$

from where we get  $\varepsilon_B^t(b_{(1)})\phi(b_{(2)}) = b_{(1)}\phi(b_{(2)})$ . Also,  $\tau(S_B(b)) = \tau(b)$  and  $\tau \circ \varepsilon_B^t(b) = \lambda^{-1}\tau(E_{M_1}(be_2)) = d^{-1}\varepsilon_B(b)$ , therefore  $\phi \circ S_B = \phi$  and  $\phi \circ \varepsilon_B^t = \varepsilon_B$ . Thus,  $\phi$  is the normalized Haar trace.

If  $H = 1$ , then the ‘trace vector’ of the restriction of  $\tau$  on  $B_t$  is given by  $\vec{\tau} = \frac{1}{d}(m_1, m_2, \dots)$ , so the componetnts of  $\vec{\tau}$  are rational numbers. Let  $\Lambda$  be the inclusion matrix of  $B_t \subset B$ , then

$$\Lambda \Lambda^t \vec{\tau} = \lambda^{-1} \vec{\tau}.$$

Since all entries of  $\Lambda \Lambda^t$  and  $\vec{\tau}$  are rational,  $\lambda^{-1}$  must be rational. On the other hand,  $\lambda^{-1}$  is an algebraic integer as an eigenvalue of the integer matrix  $\Lambda \Lambda^t$ . Therefore,  $\lambda^{-1}$  is integer.

**Proposition 4.18** *If  $N \subset M$  is a depth 2 inclusion of  $II_1$  factors such that  $[M : N]$  is a square free integer (i.e.,  $[M : N]$  is an integer which has no divisors of the form  $n^2$ ,  $n > 1$ ), then  $N' \cap M = \mathbb{C}$ , and there is a (canonical) minimal action of a Kac algebra  $B$  on  $M_1$  such that  $M_2 \cong M_1 \rtimes B$  and  $M = M_1^B$ .*

*Proof.* It suffices to show that  $N \subset M$  is irreducible, since the rest follows from [18]. Let  $q$  be a minimal projection in  $M' \cap M_1$ , then the reduced inclusion  $qM \subset qM_1q$  is of finite depth [1]. Since any finite depth inclusion is extremal (see, e.g., [16], 1.3.6) we have

$$[qM_1q : qM] = \tau(q)^2[M_1 : M] = \tau(q)^2[M : N],$$



by ([15], Corollary 4.5).

We claim that  $\tau(q)$  is a rational number. Indeed, it is well-known that the Perron-Frobenius eigenspace of the non-negative matrix  $\Lambda\Lambda^t$  is 1-dimensional [6]. Letting one of the components of a corresponding eigenvector  $\tau$  to be equal to 1, one can recover the rest of components from the system of linear equations with integer coefficients. Thus, we have that all components of  $\vec{\tau}$  are rational; clearly, the normalization condition  $\tau(1) = 1$  does not change this property.

Therefore, the index  $[qM_1q : qM]$  is a rational number. On the other hand, it must be an algebraic integer, since the depth is finite. Therefore,  $[qM_1q : qM]$  is an integer. Since  $[M : N]$  is square free, we must have  $\tau(q) = 1$ , which means that  $M' \cap M_1$  and  $N' \cap M$  are 1-dimensional.

**Corollary 4.19** *If  $N \subset M$  is a depth 2 inclusion of  $II_1$  factors such that  $[M : N] = p$  is prime, then  $N' \cap M = \mathbb{C}$ , and there is an outer action of the cyclic group  $G = \mathbb{Z}/p\mathbb{Z}$  on  $M_1$  such that  $M_2 \cong M_1 \rtimes G$  and  $M = M_1^G$ .*

*Proof.* By Proposition 4.18,  $B$  must be a Kac algebra of prime dimension  $p$ . But it is known that any such an algebra is a group algebra of the cyclic group  $G = \mathbb{Z}/p\mathbb{Z}$  [9].

## 5 Weak $C^*$ -Hopf algebra structure on $M' \cap M_2$ (the general case)

When  $H \neq 1$ ,  $(B, \Delta_B, \varepsilon_B, S_B)$  is no longer a weak Kac algebra (for instance,  $\Delta_B$  is not a homomorphism). However, it is possible to deform the structure maps in such a way that  $A$  becomes a weak  $C^*$ -Hopf algebra.

**Definition 5.1** Let us define the following operations on  $B$  :

$$\begin{aligned} \text{involution} \quad \quad \quad \dagger : B &\rightarrow B & : \quad b^\dagger &= S_B(H)^{-1}b^*S_B(H), \\ \text{comultiplication} \quad \tilde{\Delta} : B &\rightarrow B \otimes B & : \quad \tilde{\Delta}(b) &= (1 \otimes H^{-1})\Delta_B(b) \text{ i.e.,} \\ & & & b_{(\tilde{1})} \otimes b_{(\tilde{2})} = b_{(1)} \otimes H^{-1}b_{(2)} \\ \text{counit} \quad \quad \quad \tilde{\varepsilon} : B &\rightarrow \mathbb{C} & : \quad \tilde{\varepsilon}(b) &= \varepsilon_B(Hb), \\ \text{antipode} \quad \quad \quad \tilde{S} : B &\rightarrow B & : \quad \tilde{S}(b) &= S_B(HbH^{-1}). \end{aligned}$$

Clearly,  $\dagger$  defines a  $C^*$ -algebra structure on  $B$  (we will still denote this new  $C^*$ -algebra by  $B$ ). Our goal is to show that  $(B, \tilde{\Delta}, \tilde{\varepsilon}, \tilde{S})$  is a weak  $C^*$ -Hopf algebra. The proof of this fact consists of a verification of all the axioms from Section 2. We will need the following technical lemma.

**Lemma 5.2** *For all  $b \in B$  and  $z \in B_t$  we have*

- (i)  $\varepsilon_B^t(zb) = z\varepsilon_B^t(b)$ ,
- (ii)  $b_{(1)}z \otimes b_{(2)} = (bz)_{(1)} \otimes (bz)_{(2)}$ ,
- (iii)  $b_{(1)}S_B(z) \otimes b_{(2)} = b_{(1)} \otimes b_{(2)}z$ ,

*Proof.* Part (i) is clear from Proposition 4.2. Next, recall that  $B_t = A \cap B$ , and compute

$$\langle a_1, b_{(1)}z \rangle \langle a_2, b_{(2)} \rangle = \langle za_1a_2, b \rangle = \langle a_1a_2, bz \rangle, \quad a_1, a_2 \in A,$$

which gives (ii). Finally, using the properties of  $S_B$  we have

$$\begin{aligned} \langle a_1, b_{(1)}S_B(z) \rangle \langle a_2, b_{(2)} \rangle &= \langle a_1, S_B(zS_B(b_{(1)})) \rangle \langle a_2, b_{(2)} \rangle \\ &= \overline{\langle a_1^*, S_B(b_{(1)}^*)z^* \rangle} \langle a_2, b_{(2)} \rangle \\ &= \overline{\langle (a_1z)^*, S_B(b_{(1)}^*) \rangle} \langle a_2, b_{(2)} \rangle \\ &= \langle a_1z, b_{(1)} \rangle \langle a_2, b_{(2)} \rangle \\ &= \langle a_1za_2, b \rangle \\ &= \langle a_1, b_{(1)} \rangle \langle a_2, b_{(2)}z \rangle, \end{aligned}$$

from where (iii) follows.

**Proposition 5.3**  *$(B, \tilde{\Delta}, \tilde{\varepsilon})$  is a coalgebra.*

*Proof.* Let us check the coassociativity of  $\tilde{\Delta}$ . Using Lemma 5.2 and the fact that  $H \in B_t$  we compute for all  $b \in B$  :

$$\begin{aligned} (\tilde{\Delta} \otimes \text{id})\tilde{\Delta}(b) &= \tilde{\Delta}(b_{(1)}) \otimes H^{-1}b_{(2)} = b_{(1)} \otimes H^{-1}b_{(2)} \otimes H^{-1}b_{(3)} \\ &= b_{(1)} \otimes (H^{-1}b_{(2)})_{(1)} \otimes H^{-1}(H^{-1}b_{(2)})_{(2)} \\ &= b_{(1)} \otimes \tilde{\Delta}(H^{-1}b_{(2)}) = (\text{id} \otimes \tilde{\Delta})\tilde{\Delta}(b). \end{aligned}$$

Next, we check the counit axioms :

$$\begin{aligned} (\tilde{\varepsilon} \otimes \text{id})\tilde{\Delta}(b) &= \varepsilon(Hb_{(1)})H^{-1}b_{(2)} = \varepsilon((Hb)_{(1)})H^{-1}(Hb)_{(2)} = b, \\ (\text{id} \otimes \tilde{\varepsilon})\tilde{\Delta}(b) &= b_{(1)}\varepsilon(HH^{-1}b_{(2)}) = b. \end{aligned}$$

**Proposition 5.4**  $\tilde{\Delta}$  is a  $\dagger$ -homomorphism.

*Proof.* Using the properties of  $\Delta_B$  from Corollary 4.16 and Lemma 5.2 we have:

$$\begin{aligned}
\tilde{\Delta}(bc) &= (1 \otimes H^{-1})\Delta_B(bc) \\
&= (1 \otimes H^{-1})\Delta_B(b)(1 \otimes H^{-1})\Delta_B(c) = \tilde{\Delta}(b)\tilde{\Delta}(c), \\
\tilde{\Delta}(b^\dagger) &= \tilde{\Delta}(S_B(H)^{-1}b^*S_B(H)) \\
&= (S_B(H)^{-1}b^*S_B(H))_{(1)} \otimes H^{-1}(S_B(H)^{-1}b^*S_B(H))_{(2)} \\
&= S_B(H)^{-1}b_{(1)}^* \otimes S_B(H)^{-1}b_{(2)}^*S_B(H) = (S_B(H)^{-1}b_{(1)})^\dagger \otimes b_{(2)}^\dagger \\
&= b_{(1)}^\dagger \otimes (H^{-1}b_{(2)})^\dagger = \tilde{\Delta}(b)^{\dagger \otimes \dagger}.
\end{aligned}$$

**Proposition 5.5** Let  $\tilde{\varepsilon}^t(b) = \tilde{\varepsilon}(1_{(\bar{1})}b)1_{(\bar{2})}$ . Then, for all  $b, c \in B$  :

$$b\tilde{\varepsilon}^t(c) = \tilde{\varepsilon}(b_{(\bar{1})}c)b_{(\bar{2})}, \quad b_{(\bar{1})} \otimes \tilde{\varepsilon}^t(b_{(\bar{2})}) = 1_{(\bar{1})}b \otimes 1_{(\bar{2})},$$

*Proof.* First, we compute, using Lemma 5.2 and Proposition 4.3 :

$$\tilde{\varepsilon}^t(b) = \varepsilon(H1_{(1)}b)H^{-1}1_{(2)} = \varepsilon(H_{(1)}b)H_{(2)}H^{-1} = H\varepsilon_B^t(b)H^{-1} = \varepsilon_B^t(b).$$

Using this relation, Lemma 5.2, and properties of  $\varepsilon_B^t$  from Corollary 4.16 we have

$$\begin{aligned}
b_{(\bar{1})} \otimes \tilde{\varepsilon}^t(b_{(\bar{2})}) &= b_{(1)} \otimes \varepsilon_B^t(H^{-1}b_{(2)}) = b_{(1)} \otimes H^{-1}\varepsilon_B^t(b_{(2)}) \\
&= 1_{(1)}b \otimes H^{-1}1_{(2)} = 1_{(\bar{1})}b \otimes 1_{(\bar{2})}, \\
b\tilde{\varepsilon}^t(c) &= b\varepsilon_B^t(c) = H^{-1}(Hb)\varepsilon_B^t(c) \\
&= H^{-1}\varepsilon_B((Hb)_{(1)}c)(Hb)_{(2)} = \varepsilon_B(Hb_{(1)}c)H^{-1}b_{(2)} \\
&= \tilde{\varepsilon}(b_{(\bar{1})}c)b_{(\bar{2})}.
\end{aligned}$$

**Proposition 5.6**  $\tilde{S}$  is a linear anti-multiplicative and anti-comultiplicative map such that

$$b_{(\bar{1})}\tilde{S}(b_{(\bar{2})}) = \tilde{\varepsilon}^t(b).$$

Moreover,  $(\tilde{S} \circ \dagger)^2 = id$  and  $\tilde{S}^2(b) = GbG^{-1}$ , where  $G = \tilde{S}(H)^{-1}H$ .

*Proof.* Using Corollary 4.16, Lemma 5.2 and definitions of  $\tilde{S}$  and  $\dagger$ , we have :

$$\begin{aligned}
\tilde{S}(bc) &= S_B(HbcH^{-1}) = S_B(HcH^{-1})S_B(HbH^{-1}) = \tilde{S}(c)\tilde{S}(b), \\
\tilde{S}(b)_{(\tilde{2})} \otimes \tilde{S}(b)_{(\tilde{1})} &= H^{-1}S_B(HbH^{-1})_{(2)} \otimes S_B(HbH^{-1})_{(1)} \\
&= S_B(HbH^{-1})_{(2)} \otimes S_B(H^{-1})S_B(HbH^{-1})_{(1)} \\
&= S_B((HbH^{-1})_{(1)}) \otimes S_B(H^{-1})S_B((HbH^{-1})_{(2)}) \\
&= S_B(Hb_{(1)}H^{-1}) \otimes S_B(b_{(2)}H^{-1}) \\
&= \tilde{S}(b_{(1)}) \otimes \tilde{S}(H^{-1}b_{(2)}) = \tilde{S}(b_{(\tilde{1})}) \otimes \tilde{S}(b_{(\tilde{2})}), \\
b_{(\tilde{1})}\tilde{S}(b_{(\tilde{2})}) &= b_{(1)}S_B(b_{(2)}H^{-1}) = \varepsilon_B^t(b) = \tilde{\varepsilon}^t(b),
\end{aligned}$$

from where the first part of Proposition follows. Next, we can compute

$$\begin{aligned}
\tilde{S}(b^\dagger) &= S_B(Hb^\dagger H^{-1}) = S_B(HS_B(H)^{-1}b^*S_B(H)H^{-1}) \\
&= S_B(H)^{-1}HS_B(b^*)H^{-1}S_B(H), \\
\tilde{S}(\tilde{S}(b^\dagger)^\dagger) &= \tilde{S}((S_B(H)^{-1}HS_B(b^*)H^{-1}S_B(H))^\dagger) = \tilde{S}(H^{-1}S_B(b)H) \\
&= S_B(S_B(b)) = b,
\end{aligned}$$

therefore  $(\tilde{S} \circ \dagger)^2 = \text{id}$ . Finally, since  $S_B(H) = \tilde{S}(H)$ , we get

$$\begin{aligned}
\tilde{S}^2(b) &= \tilde{S}(S_B(HbH^{-1})) = S_B(HS_B(HbH^{-1})H^{-1}) \\
&= S_B(H)^{-1}HbH^{-1}S_B(H) = GbG^{-1}.
\end{aligned}$$

Thus, we can state the main result of this section.

**Theorem 5.7** *( $B, \tilde{\Delta}, \tilde{\varepsilon}, \tilde{S}$ ) is a weak  $C^*$ -Hopf algebra with the Haar projection  $e_2H$  and normalized Haar functional  $\tilde{\phi}(b) = \phi(H\tilde{S}(H)b) = d\tau(\tilde{S}(H)Hb)$  (cf. Theorem 4.17).*

*Proof.* It follows from Propositions 5.3 – 5.6 that  $(B, \tilde{\Delta}, \tilde{\varepsilon}, \tilde{S})$  is a weak  $C^*$ -Hopf algebra.

The properties of  $e_2$  established in Theorem 4.17 and Proposition 4.2 give

$$\begin{aligned}
be_2H &= \varepsilon_B^t(b)e_2H = \tilde{\varepsilon}^t(b)e_2H, \\
\tilde{\varepsilon}^t(e_2H) &= \varepsilon_B^t(e_2H) = \lambda^{-1}E_{M_1}(e_2He_2) \\
&= \lambda^{-1}E_{M_1}(E_M(H)e_2) = E_M(H) = 1,
\end{aligned}$$

since  $\tilde{\varepsilon}^t = \varepsilon_B^t$  by Proposition 5.5, and  $E_M(H) = \tau(H)1 = 1$  by Corollary 4.7.

Using Lemma 5.2(ii), and taking into account that  $\tilde{S}^2|_{B_t} = \text{id}_{B_t}$  (Proposition 5.6), we compute for all  $b \in B$ ,  $z \in B_t$  :

$$\tilde{\varepsilon}^t(\tilde{S}(z)) = \tilde{S}(z)_{(\tilde{1})} \tilde{S}(\tilde{S}(z)_{(\tilde{2})}) = 1_{(\tilde{1})} \tilde{S}(\tilde{S}(z)1_{(\tilde{2})}) = \tilde{S}^2(z) = z,$$

therefore  $e_2 \tilde{S}(H) = e_2 \tilde{\varepsilon}^t(\tilde{S}(H))^* = e_2 H$ . Since  $S_B(e_2) = e_2$  and  $S_B(H) = \tilde{S}(H)$ , using the above relation, we get

$$\tilde{S}(e_2 H) = \tilde{S}(H)^{-1} S_B(e_2 H) \tilde{S}(H) = e_2 \tilde{S}(H) = e_2 H.$$

Thus  $\tilde{S}(e_2 H) = e_2 H$ . Also we have :

$$\begin{aligned} (e_2 H)^2 &= E_M(H) e_2 H = e_2 H, \\ (e_2 H)^\dagger &= \tilde{S}(H)^{-1} H e_2 \tilde{S}(H) = e_2 \tilde{S}(H) = e_2 H. \end{aligned}$$

Therefore,  $e_2 H$  is an  $\tilde{S}$ -invariant projection. This proves that  $e_2$  is the Haar projection of  $B$ .

Next, using Lemma 5.2 and the properties of the trace  $\phi$  from the proof of Theorem 4.17 we have

$$\begin{aligned} \tilde{\varepsilon}^t(b_{(\tilde{1})}) \tilde{\phi}(b_{(\tilde{2})}) &= \varepsilon_B^t(b_{(1)}) \tilde{\phi}(H^{-1}b_{(2)}) = \varepsilon_B^t(b_{(1)}) \phi(\tilde{S}(H)b_{(2)}) \\ &= \varepsilon_B^t((b\tilde{S}(H))_{(1)}) \phi((b\tilde{S}(H))_{(2)}) = (b\tilde{S}(H))_{(1)} \phi((b\tilde{S}(H))_{(2)}) \\ &= b_{(1)} \phi(\tilde{S}(H)b_{(2)}) = b_{(1)} \phi(H\tilde{S}(H)H^{-1}b_{(2)}) \\ &= b_{(1)} \tilde{\phi}(H^{-1}b_{(2)}) = b_{(\tilde{1})} \tilde{\phi}(b_{(\tilde{2})}), \\ \tilde{\phi}(\tilde{S}(b)) &= \phi(H\tilde{S}(H)\tilde{S}(H)^{-1}S_B(b)\tilde{S}(H)) = \phi(H\tilde{S}(H)S_B(b)) \\ &= \phi(b\tilde{S}(H)H) = \tilde{\phi}(b), \\ \tilde{\phi}(\tilde{\varepsilon}^t(b)) &= \phi(\tilde{S}(H)H\varepsilon_B^t(b)) = \tau(\tilde{S}(H))\phi(\varepsilon_B^t(Hb)) = \varepsilon_B(Hb) = \tilde{\varepsilon}(b), \end{aligned}$$

therefore,  $\tilde{\phi}$  is the normalized Haar functional on  $B$ .

**Remark 5.8** (i) The non-degenerate duality  $<, >$  induces on  $A = N' \cap M_1$  the structure of the weak  $C^*$ -Hopf algebra dual to  $B$ .

(ii) The weak  $C^*$ -Hopf algebra  $B$  is biconnected, since the inclusion  $B_t = M' \cap M_1 \subset B = M' \cap M_2$  is connected ([7], 4.6.3) and  $B_t \cap B_s = (M' \cap M_1) \cap (M'_1 \cap M_2) = \mathbb{C}$ . Thus, only biconnected weak Hopf  $C^*$ -algebras arise as symmetries of finite index depth 2 inclusions of  $\text{II}_1$  factors.

- (iii) If  $\lambda^{-1}$  is not integer, then  $\tilde{S}$  has infinite order. Indeed, the canonical element  $G$  implementing the square of the antipode in Proposition 5.6 is positive, so if  $\tilde{S}^{2n} = \text{id}$  for some  $n$ , then  $G^n$  belongs to  $Z(B)$ , the center of  $B$ . Taking the  $n$ -th root, we get that  $G \in Z(B)$ , which means that  $S^2 = \text{id}$ , and  $B$  is a weak Kac algebra, which is in contradiction with Theorem 4.17.

## 6 Action of $B$ on $M_1$ .

Note that in terms of  $\tilde{\Delta}$ , Proposition 4.13 means that

$$E_{M_1}(bx ye_2) = \lambda^{-1} E_{M_1}(b_{(\tilde{1})} x e_2) E_{M_1}(b_{(\tilde{2})} y e_2),$$

for all  $b \in B$  and  $x, y \in M_1$ . This suggests the following definition of the action of  $B$  on  $M_1$ .

**Proposition 6.1** *The map  $\triangleright : B \otimes M_1 \rightarrow M_1$  :*

$$b \triangleright x = \lambda^{-1} E_{M_1}(b x e_2)$$

*defines a left action of  $B$  on  $M_1$  (cf. [18], Proposition 17).*

*Proof.* Clearly, the above map defines a left  $B$ -module structure on  $M_1$ , since  $1 \triangleright x = x$  and

$$b \triangleright (c \triangleright x) = \lambda^{-2} E_{M_1}(b E_{M_1}(c x e_2) e_2) = \lambda^{-1} E_{M_1}(b c x e_2) = (bc) \triangleright x.$$

Next, using Proposition 4.13 we get

$$\begin{aligned} b \triangleright xy &= \lambda^{-1} E_{M_1}(b x y e_2) = \lambda^{-2} E_{M_1}(b_{(\tilde{1})} x e_2) E_{M_1}(b_{(\tilde{2})} y e_2) \\ &= (b_{(\tilde{1})} \triangleright x) (b_{(\tilde{2})} \triangleright y). \end{aligned}$$

By Remark 4.4 and properties of  $S_B$  we also get

$$\begin{aligned} \tilde{S}(b)^\dagger \triangleright x^* &= \lambda^{-1} E_{M_1}(\tilde{S}(b)^\dagger x^* e_2) \\ &= \lambda^{-1} E_{M_1}(S_B(H)^{-1} S_B(H b H^{-1})^* S_B(H) x^* e_2) \\ &= \lambda^{-1} E_{M_1}(S_B(b^*) x^* e_2) \\ &= \lambda^{-1} E_{M_1}(e_2 x^* b^*) = \lambda^{-1} E_{M_1}(b x e_2)^* = (b \triangleright x)^*. \end{aligned}$$

Finally,

$$b \triangleright 1 = \lambda^{-1} E_{M_1}(b e_2) = \lambda^{-1} E_{M_1}(\lambda^{-1} E_{M_1}(b e_2) e_2) = \tilde{\varepsilon}^t(b) \triangleright 1,$$

and  $b \triangleright 1 = 0$  iff  $\tilde{\varepsilon}^t(b) = \lambda^{-1} E_{M_1}(b e_2) = 0$ .

**Proposition 6.2**  $M_1^B = M$ , i.e.  $M$  is the fixed point subalgebra of  $M_1$ .

*Proof.* If  $x \in M_1$  is such that  $b \triangleright x = \tilde{\varepsilon}^t(b) \triangleright x$  for all  $b \in B$ , then  $E_{M_1}(bx e_2) = E_{M_1}(\varepsilon_B^t(b)x e_2) = E_{M_1}(b e_2)x$ . Taking  $b = e_2$ , we get  $E_M(x) = x$  which means that  $x \in M$ . Thus,  $M_1^B \subset M$ .

Conversely, if  $x \in M$ , then  $x$  commutes with  $e_2$  and

$$b \triangleright x = \lambda^{-1} E_{M_1}(b e_2 x) = \lambda^{-1} E_{M_1}(\lambda^{-1} E_{M_1}(b e_2) e_2 x) = \varepsilon_B^t(b) \triangleright x,$$

therefore  $M_1^B = M$ .

**Proposition 6.3** The map  $\theta : [x \otimes b] \mapsto x \tilde{S}(H)^{\frac{1}{2}} b \tilde{S}(H)^{-\frac{1}{2}}$  defines a von Neumann algebra isomorphism between  $M_1 \rtimes B$  and  $M_2$ .

*Proof.* By definition of the action  $\triangleright$  we have :

$$\begin{aligned} \theta([x(z \triangleright 1) \otimes b]) &= x \tilde{S}(H)^{\frac{1}{2}} \lambda^{-1} E_{M_1}(z e_2) b \tilde{S}(H)^{-\frac{1}{2}} \\ &= x \tilde{S}(H)^{\frac{1}{2}} z b \tilde{S}(H)^{-\frac{1}{2}} = \theta([x \otimes z b]), \end{aligned}$$

for all  $x \in M_1$ ,  $b \in B$ ,  $z \in B_t$ , so  $\theta$  is a well defined linear map from  $M_1 \rtimes B = M_1 \otimes_{B_t} B$  to  $M_2$ . It is surjective since an orthonormal basis of  $B = M' \cap M_2$  over  $B_t = M' \cap M_1$  is also a basis of  $M_2$  over  $M_1$  ([16], 2.1.3).

Let us check that  $\theta$  is an involution-preserving isomorphism of algebras. Note that from Corollary 4.12 we have  $bx = (b_{(\bar{1})} \triangleright x) b_{(\bar{2})}$ . This allows us to compute, for all  $x, y \in M_1$  and  $b, c \in B$ :

$$\begin{aligned} \theta([x \otimes b][y \otimes c]) &= \theta([x(b_{(\bar{1})} \triangleright y) \otimes b_{(\bar{2})} c]) \\ &= x(b_{(\bar{1})} \triangleright y) \tilde{S}(H)^{\frac{1}{2}} b_{(\bar{2})} c \tilde{S}(H)^{-\frac{1}{2}} \\ &= x((\tilde{S}(H)^{\frac{1}{2}} b)_{(\bar{1})} \triangleright y) (\tilde{S}(H)^{\frac{1}{2}} b)_{(\bar{2})} c \tilde{S}(H)^{-\frac{1}{2}} \\ &= x \tilde{S}(H)^{\frac{1}{2}} b y c \tilde{S}(H)^{-\frac{1}{2}} \\ &= x \tilde{S}(H)^{\frac{1}{2}} b \tilde{S}(H)^{-\frac{1}{2}} y \tilde{S}(H)^{\frac{1}{2}} c \tilde{S}(H)^{-\frac{1}{2}} \\ &= \theta([x \otimes b]) \theta([y \otimes c]), \\ \theta([x \otimes b]^*) &= (b_{(\bar{1})}^\dagger \triangleright x^*) \tilde{S}(H)^{\frac{1}{2}} b_{(\bar{2})}^\dagger \tilde{S}(H)^{-\frac{1}{2}} \\ &= (\tilde{S}(H)^{\frac{1}{2}} b^\dagger)_{(\bar{1})} \triangleright x^*) (\tilde{S}(H)^{\frac{1}{2}} b^\dagger)_{(\bar{2})} \tilde{S}(H)^{-\frac{1}{2}} \\ &= \tilde{S}(H)^{\frac{1}{2}} b^\dagger \tilde{S}(H)^{-\frac{1}{2}} x^* \\ &= \tilde{S}(H)^{-\frac{1}{2}} b^* \tilde{S}(H)^{\frac{1}{2}} x^* \\ &= (x \tilde{S}(H)^{\frac{1}{2}} b \tilde{S}(H)^{-\frac{1}{2}})^* = \theta([x \otimes b])^*. \end{aligned}$$

It is known that  $M_1 \rtimes B$  is a  $\text{II}_1$  factor iff  $M_1^B$  is [13]. Now the injectivity of  $\theta$  follows from the simplicity of  $\text{II}_1$  factors (see, e.g., the appendix of [8]). Thus,  $\theta$  is a von Neumann algebra isomorphism.

- Remark 6.4** (i) The action of  $B$  constructed in Proposition 6.1 is minimal, since we have  $M_1' \cap M_1 \rtimes B = M_1' \cap M_2 = B_s$  by Proposition 6.3.
- (ii) If the inclusion  $N \subset M$  is irreducible, then  $B$  is a usual Kac algebra (i.e., a Hopf  $C^*$ -algebra) and we recover the well-known result proved in [18], [10], and [3].

## References

- [1] D. Bisch. On the structure of finite depth subfactors. In : Algebraic methods in Operator Theory, Birkhäuser, (1994) 175–194.
- [2] G. Böhm, F. Nill and K. Szlachányi. Weak Hopf algebras I: Integral theory and  $C^*$ -structure. *math.QA/9805116* (1998).
- [3] M.-C. David. Paragroupe d’Adrian Ocneanu et algebre de Kac. *Pacif. J. of Math.*, **172**, no. 2 (1996), 331–363.
- [4] M. Enock and R. Nest. Inclusions of factors, multiplicative unitaries and Kac Algebras. *J. Funct. Analysis*, **137** (1996), 466–543.
- [5] M. Enock and J.-M. Vallin. Inclusions of von Neumann Algebras and quantum groupoids. *Inst. Math. de Jussieu*, Prépublication 156 (1998).
- [6] F.R. Gantmacher. Applications of the theory of matrices, Interscience Publishers, (1959).
- [7] F.M. Goodman, P. de la Harpe, and V.F.R. Jones. Coxeter graphs and towers of algebras. MSRI Publ. 14, Springer-Verlag, (1989).
- [8] V. Jones and V.S. Sunder. Introduction to subfactors, Cambridge University Press, (1997).
- [9] G.I. Kac. Certain arithmetic properties of ring groups. *Functional Anal. Appl.*, **6** (1972), 158–160.
- [10] R. Longo. A duality for Hopf algebras and subfactors I. *Comm. Math. Phys.*, **159** (1994), 133–150.



- [11] D. Nikshych, L. Vainerman. Algebraic versions of a finite dimensional quantum groupoid. *math.QA/9808054* (1998).
- [12] D. Nikshych. Duality for actions of weak Kac algebras and crossed product inclusions of  $\text{II}_1$  factors. *math.QA/9810049* (1998).
- [13] F. Nill, K. Szlachányi, H.-W. Wiesbrock. Weak Hopf algebras and reducible Jones inclusions of depth 2. I: From crossed products to Jones towers. *math.QA/9806130* (1998).
- [14] F. Nill. Axioms for weak bialgebras. *math.QA/9805104* (1998)
- [15] M. Pimsner and S. Popa. Entropy and index for subfactors. *Ann. Sci. Ecole Norm. Sup. (4)* **19** (1986), no. 3, 57–106.
- [16] S. Popa. Classification of amenable subfactors of type II. *Acta Math.* **172** (1994), 163–255.
- [17] S. Popa. Classification of subfactors : the reduction to the commuting squares. *Invent. Math.* **101** (1990), 19–43.
- [18] W. Szymański. Finite index subfactors and Hopf algebra crossed products. *Proc. Amer. Math. Soc.* **120** (1994), no. 2, 519–528.
- [19] J.-M. Vallin. Bimodules de Hopf et poids opératoriels de Haar, *J. Operator Theory*, **35** (1996), 39–65.
- [20] Y. Watatani. Index for  $C^*$ -subalgebras, *Memoirs of the AMS* **424** (1990).
- [21] T. Yamanouchi. Duality for generalized Kac algebras and a characterization of finite groupoid algebras. *J. Algebra*, **163** (1994), 9–50.